


# SUSY naturalness, PQ symmetry and the landscape



Howard Baer  
University of Oklahoma

Is SUSY Alive and Well?  
Madrid, Sept. 2016

twin pillars of guidance:  
naturalness & simplicity



“The appearance of fine-tuning in a scientific theory is like a cry of distress from nature, complaining that something needs to be better explained”



S. Weinberg

“Everything should be made as simple as possible, but not simpler”


A. Einstein

# Nature sure looks like SUSY

- stabilize Higgs mass
- measured gauge couplings
- $m(t) \sim 173$  GeV for REWSB
- $m_h(125)$ : squarely within SUSY window



recent search results from Atlas run 2 @ 13 TeV:




evidently  $m_{\tilde{g}} > 1.9 \text{ TeV}$

compare: BG naturalness (1987):  $m_{\tilde{g}} < 0.35 \text{ TeV}$

These bounds appear in sharp conflict with EW “naturalness”

	mass
gluino	400 GeV
uR	400 GeV
eR	350 GeV
chargino	100 GeV
neutralino	50 GeV



Cassel, Ghilencea, Ross, 2009

$\Delta \rightarrow 1000$   
as  $m_h \rightarrow 125$  GeV  
0.1% tuning!?

Barbieri-Giudice 10% bounds, 1987

# IS SUSY ALIVE AND WELL?




Instituto de Física Teórica UAM-CSIC  
Madrid, 28-30 September 2016

<https://workshops.ift.uam-csic.es/susyaaw>

or is SUSY dead?  
how to disprove SUSY?  
when it becomes “unnatural”?  
this brings up **naturalness issue**

Mark Twain, 1835-1910 (or SUSY)



1897

“...settling the ultimate fate of naturalness is perhaps the most profound theoretical question of our time”



Arkani-Hamed et al.,  
arXiv:1511.06495

“Given the magnitude of the stakes involved,  
it is vital to get a clear verdict  
on naturalness from experiment”

This should be matched by theoretical scrutiny  
of what we mean by naturalness

# Oft-repeated **myths** about naturalness

- requires  $m(t_1, t_2, b_1) < 500$  GeV
- requires small  $A_t$  parameter
- requires  $m(\text{gluino}) < 400$ , no 1000, no 1500 GeV?
- MSSM is fine-tuned to .1% - needs modification
- naturalness is subjective/ non-predictive
- different measures predict different things

**This talk will refute all these points!**

HB, Barger, Savoy, arXiv:1502.04127

**And present a beautiful alternative:  
radiatively-driven naturalness**



Most claims against SUSY stem from **overestimates** of EW fine-tuning.

These arise from violations of the

**Prime directive on fine-tuning:**

“Thou shalt not claim fine-tuning of **dependent** quantities one against another!”

HB, Barger, Mickelson, Padeffke-Kirkland, arXiv:1404.2277




Is  $\mathcal{O} = \mathcal{O} + b - b$  fine-tuned for  $b > \mathcal{O}$ ?

## Reminder: naturalness in the SM

Higgs sector of SM is “natural” only up to cutoff

$$\begin{aligned} V &= -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 \\ m_h^2 &\simeq 2\mu^2 + \delta m_h^2 \\ \delta m_h^2 &\simeq \frac{3}{4\pi^2} \left( -\lambda_t^2 + \frac{g^2}{4} + \frac{g^2}{8 \cos^2 \theta_W} + \lambda \right) \Lambda^2 \end{aligned}$$




Since  $\delta m_h^2$  is *independent* of  $\mu^2$ ,  
can freely dial (fine-tune)  $\mu^2$  to maintain  $m_h = 125$  GeV


Naturalness:  $\delta m_h^2 < m_h^2 \Rightarrow \Lambda < 1$  TeV!  
New physics at or around the TeV scale!

Next: simple electroweak fine-tuning in SUSY:  
 dial value of  $\mu$  so that Z mass comes out right:  
 everybody does it but it is hidden inside spectra  
 codes (Isajet, SuSpect, SoftSUSY, Spheno, SSARD)


$$\frac{m_Z^2}{2} = \frac{m_{H_d}^2 + \Sigma_d^d - (m_{H_u}^2 + \Sigma_u^u) \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2 \simeq -m_{H_u}^2 - \Sigma_u^u - \mu^2$$



e.g. in CMSSM/  
 mSUGRA:  
 one then concludes  
 nature  
 gives this:



If you didn't fine-tune, then here is  $m(Z)$



The 20 dimensional pMSSM parameter space then includes

$M_1, M_2, M_3,$   
 $m_{Q_1}, m_{U_1}, m_{D_1}, m_{L_1}, m_{E_1},$   
 $m_{Q_3}, m_{U_3}, m_{D_3}, m_{L_3}, m_{E_3},$   
 $A_t, A_b, A_\tau,$   
 $m_{H_u}^2, m_{H_d}^2, \mu, B.$

scan over parameters

Natural value of  $m(Z)$  from  
pMSSM is  $\sim 2-4$  TeV

# Three measures of fine-tuning:



# #1: Simplest SUSY measure: $\Delta_{EW}$

Working only at the weak scale, minimize scalar potential: calculate  $m(Z)$  or  $m(h)$

No large uncorrelated cancellations in  $m(Z)$  or  $m(h)$

$$\frac{m_Z^2}{2} = \frac{m_{H_d}^2 + \Sigma_d^d - (m_{H_u}^2 + \Sigma_u^u) \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2 \sim -m_{H_u}^2 - \Sigma_u^u - \mu^2$$

$$\Delta_{EW} \equiv \max_i |C_i| / (m_Z^2/2) \quad \text{with} \quad C_{H_u} = -m_{H_u}^2 \tan^2 \beta / (\tan^2 \beta - 1) \quad \text{etc.}$$

simple, direct, unambiguous interpretation:

- $|\mu| \sim m_Z \sim 100 - 200 \text{ GeV}$
- $m_{H_u}^2$  should be driven to small negative values such that  $-m_{H_u}^2 \sim 100 - 200 \text{ GeV}$  at the weak scale and
- that the radiative corrections are not too large:  $\Sigma_u^u \lesssim 100 - 200 \text{ GeV}$

CETUP\*-12/002, FTPI-MINN-12/22, UMN-TH-3109/12, UH-511-1195-12

Radiative natural SUSY with a 125 GeV Higgs boson

Howard Baer,<sup>1</sup> Vernon Barger, Peisi Huang,<sup>2</sup> Azar Mustafayev,<sup>3</sup> and Xerxes Tata<sup>4</sup>


<sup>1</sup>Dept. of Physics and Astronomy, University of Oklahoma, Norman, OK, 73019, USA

<sup>2</sup>Dept. of Physics, University of Wisconsin, Madison, WI 53706, USA

<sup>3</sup>W. I. Fine Institute for Theoretical Physics, University of Minnesota, Minneapolis, MN 55455, USA



PRL109 (2012) 161802

radiative corrections drive  $m_{H_u}^2$  from unnatural GUT scale values to naturalness at weak scale:  
radiatively-driven naturalness



Evolution of the soft SUSY breaking mass squared term  $sign(m_{H_u}^2)\sqrt{|m_{H_u}^2|}$  vs.  $Q$

Large value of  $A_t$  reduces  $\Sigma_u^u(\tilde{t}_{1,2})$  contributions to  $\Delta_{EW}$  while uplifting  $m_h$  to  $\sim 125$  GeV



$$\Sigma_u^u(\tilde{t}_{1,2}) = \frac{3}{16\pi^2} F(m_{\tilde{t}_{1,2}}^2) \left[ f_t^2 - g_Z^2 \mp \frac{f_t^2 A_t^2 - 8g_Z^2 (\frac{1}{4} - \frac{2}{3}x_W) \Delta_t}{m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2} \right]$$

$$\Delta_t = (m_{\tilde{t}_L}^2 - m_{\tilde{t}_R}^2)/2 + M_Z^2 \cos 2\beta (\frac{1}{4} - \frac{2}{3}x_W)$$

$$F(m^2) = m^2 \left( \log \frac{m^2}{Q^2} - 1 \right) \quad Q^2 = m_{\tilde{t}_1} m_{\tilde{t}_2}$$



## #2: Higgs mass or large-log fine-tuning $\Delta_{HS}$

It is tempting to pick out one-by-one quantum fluctuations **but** must combine log divergences before taking any limit

$$m_h^2 \simeq \mu^2 + m_{H_u}^2 + \delta m_{H_u}^2|_{rad}$$

$$\frac{dm_{H_u}^2}{dt} = \frac{1}{8\pi^2} \left( -\frac{3}{5}g_1^2 M_1^2 - 3g_2^2 M_2^2 + \frac{3}{10}g_1^2 S + 3f_t^2 X_t \right) \quad X_t = m_{Q_3}^2 + m_{U_3}^2 + m_{H_u}^2 + A_t^2$$

neglect gauge pieces, S,  $m_{H_u}$  and running;  
then we can integrate from  $m(SUSY)$  to  $\Lambda$

$$\delta m_{H_u}^2 \sim -\frac{3f_t^2}{8\pi^2} (m_{Q_3}^2 + m_{U_3}^2 + A_t^2) \ln(\Lambda/m_{SUSY})$$

$$\Delta_{HS} \sim \delta m_h^2 / (m_h^2/2) < 10$$

$$m_{\tilde{t}_{1,2}, \tilde{b}_1} < 500 \text{ GeV}$$

$$m_{\tilde{g}} < 1.5 \text{ TeV}$$

**old natural SUSY**


then

$A_t$  can't be too big

What's wrong with this argument?  
 In zeal for simplicity, have made several simplifications: most **egregious** is that one sets  $m(H_u)^2=0$  at beginning to simplify

$m_{H_u}^2(\Lambda)$  and  $\delta m_{H_u}^2$  are *not* independent!

**violates prime directive!**



The larger  $m_{H_u}^2(\Lambda)$  becomes, then the larger becomes the cancelling correction!

HB, Barger, Savoy

To fix: combine dependent terms:

$$m_h^2 \simeq \mu^2 + (m_{H_u}^2(\Lambda) + \delta m_{H_u}^2) \text{ where now both } \mu^2 \text{ and } (m_{H_u}^2(\Lambda) + \delta m_{H_u}^2) \text{ are } \sim m_Z^2$$

After re-grouping:  $\Delta_{HS} \simeq \Delta_{EW}$

Instead of: the radiative correction  $\delta m_{H_u}^2 \sim m_Z^2$   
we now have: the radiatively-corrected  $m_{H_u}^2 \sim m_Z^2$

Recommendation: put this horse out to pasture

$$\delta m_{H_u}^2 \sim -\frac{3f_t^2}{8\pi^2} (m_{Q_3}^2 + m_{U_3}^2 + A_t^2) \ln(\Lambda/m_{SUSY})$$

R.I.P.

sub-TeV 3rd generation squarks **not** required for naturalness

### #3. What about EENZ/BG measure?

$$\Delta_{BG} = \max_i \left| \frac{\partial \log m_Z^2}{\partial \log p_i} \right| = \max_i \left| \frac{p_i}{m_Z^2} \frac{\partial m_Z^2}{\partial p_i} \right|$$

applied to pMSSM, then  $\Delta_{BG} \simeq \Delta_{EW}$

What if we apply to high (e.g. GUT) scale parameters ?

$$\begin{aligned} m_Z^2 \simeq & -2.18\mu^2 + 3.84M_3^2 + 0.32M_3M_2 + 0.047M_1M_3 - 0.42M_2^2 \\ & + 0.011M_2M_1 - 0.012M_1^2 - 0.65M_3A_t - 0.15M_2A_t \\ & - 0.025M_1A_t + 0.22A_t^2 + 0.004M_3A_b \\ & - 1.27m_{H_u}^2 - 0.053m_{H_d}^2 \\ & \hline & + 0.73m_{Q_3}^2 + 0.57m_{U_3}^2 + 0.049m_{D_3}^2 - 0.052m_{L_3}^2 + 0.053m_{E_3}^2 \\ & \hline & + 0.051m_{Q_2}^2 - 0.11m_{U_2}^2 + 0.051m_{D_2}^2 - 0.052m_{L_2}^2 + 0.053m_{E_2}^2 \\ & \hline & + 0.051m_{Q_1}^2 - 0.11m_{U_1}^2 + 0.051m_{D_1}^2 - 0.052m_{L_1}^2 + 0.053m_{E_1}^2, \end{aligned}$$

For correlated scalar masses  $\equiv m_0$ ,

scalar contribution collapses:

what looks fine-tuned isn't: *focus point SUSY*

multi-TeV scalars are *natural*

Feng, Matchev, Moroi

What about EENZ/BG measure?

$$\Delta_{BG} = \max_i \left| \frac{\partial \log m_Z^2}{\partial \log p_i} \right| = \max_i \left| \frac{p_i}{m_Z^2} \frac{\partial m_Z^2}{\partial p_i} \right|$$

applied to pMSSM, then  $\Delta_{BG} \simeq \Delta_{EW}$

apply to high (e.g. GUT) scale parameters

$$\begin{aligned} m_Z^2 \simeq & -2.18\mu^2 + 3.84M_3^2 + 0.32M_3M_2 + 0.047M_1M_3 - 0.42M_2^2 \\ & + 0.011M_2M_1 - 0.012M_1^2 - 0.65M_3A_t - 0.15M_2A_t \\ & - 0.025M_1A_t + 0.22A_t^2 + 0.004M_3A_b \\ & - 1.27m_{H_u}^2 - 0.053m_{H_d}^2 \\ & + 0.73m_{Q_3}^2 + 0.57m_{U_3}^2 + 0.049m_{D_3}^2 - 0.052m_{L_3}^2 + 0.053m_{E_3}^2 \\ & + 0.051m_{Q_2}^2 - 0.11m_{U_2}^2 + 0.051m_{D_2}^2 - 0.052m_{L_2}^2 + 0.053m_{E_2}^2 \\ & + 0.051m_{Q_1}^2 - 0.11m_{U_1}^2 + 0.051m_{D_1}^2 - 0.052m_{L_1}^2 + 0.053m_{E_1}^2, \end{aligned}$$

applied to most parameters,

$\Delta_{BG}$  large, looks fine-tuned for *e.g.*  $m_{\tilde{g}} \simeq M_3 > 1.8 \text{ TeV}$

$$\Delta_{BG}(M_3^2) = 3.84 \frac{M_3^2}{m_z^2} \simeq 1500$$

But wait! in more complete models,  
soft terms not independent

violates prime directive!

e.g. in SUGRA, for well-specified hidden sector,  
each soft term calculated as multiple of  $m_{3/2}$ ;  
soft terms must be combined!

e.g. dilaton-dominated SUSY breaking:  $m_0^2 = m_{3/2}^2$  with  $m_{1/2} = -A_0 = \sqrt{3}m_{3/2}$

$$m_{H_u}^2 = a_{H_u} \cdot m_{3/2}^2,$$

$$m_{Q_3}^2 = a_{Q_3} \cdot m_{3/2}^2,$$

$$A_t = a_{A_t} \cdot m_{3/2},$$

$$M_i = a_i \cdot m_{3/2},$$

....


since  $\mu$  hardly runs, then

$$\begin{aligned} m_Z^2 &\simeq -2\mu^2 + a \cdot m_{3/2}^2 \\ &\simeq -2\mu^2 - 2m_{H_u}^2 (weak) \end{aligned}$$

$$m_{H_u}^2 (weak) \sim -(100 - 200)^2 \text{ GeV}^2 \sim -a \cdot m_{3/2}^2/2$$

using  $\mu^2$  and  $m_{3/2}^2$  as fundamental,  
then  $\Delta_{BG} \simeq \Delta_{EW}$  even using high scale parameters!

# How much is too much fine-tuning?




Visually, large fine-tuning has already developed by  $\mu \sim 350$  or  $\Delta_{EW} \sim 30$




$\Delta_{EW}$  is highly selective:  
 most constrained models are ruled out  
 except NUHM2 and its generalizations:

J. Ellis, K. Olive and Y. Santoso, *Phys. Lett. B* 539 (2002) 107; J. Ellis, T. Falk, K. Olive and Y. Santoso, *Nucl. Phys. B* 652 (2003) 259; H. Baer, A. Mustafayev, S. Profumo, A. Belyaev and X. Tata, *J. High Energy Phys.* 0507 (2005) 065.

scan over p-space with  $m(h)=125.5\pm 2.5$  GeV:



Applied properly, all three measures agree:  
**naturalness is unambiguous and highly predictive!**



Radiatively-driven natural SUSY, or RNS:

(typically need  $m_{H_u} \sim 25\text{-}50\%$  higher than  $m_0$ )


H. Baer, V. Barger, P. Huang, A. Mustafayev and X. Tata, *Phys. Rev. Lett.* **109** (2012) 161802.

H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, *Phys. Rev. D* **87** (2013) 115028 [arXiv:1212.2655 [hep-ph]].

bounds from naturalness (3%)	BG/DG	Delta_EW
mu	350 GeV	350 GeV
gluino	400-600 GeV	4000 GeV
t1	450 GeV	3000 GeV
sq/sl	550-700 GeV	10-20 TeV

h(125) and LHC limits are perfectly compatible with 3-10% naturalness: **no crisis!**

# Typical spectrum for low $\Delta_{EW}$ models



There is a Little Hierarchy, but it is **no problem**

$$\mu \ll m_{3/2}$$

SUSY  $\mu$  problem:  $\mu$  term is SUSY, not SUSY breaking:  
expect  $\mu \sim M(\text{Pl})$  but phenomenology requires  $\mu \sim m(\text{Z})$

- NMSSM:  $\mu \sim m(3/2)$ ; beware singlets!
- Giudice–Masiero:  $\mu$  forbidden by some symmetry:  
generate via Higgs coupling to hidden sector
- **Kim–Nilles**: invoke SUSY version of DFSZ axion  
solution to strong CP:

KN: PQ symmetry forbids  $\mu$  term,  
but then it is generated via PQ breaking

Little Hierarchy due to mismatch between  
PQ breaking and SUSY breaking scales?

$$\mu \sim \lambda f_a^2 / M_P$$

$$m_{3/2} \sim m_{hid}^2 / M_P$$

$$f_a \ll m_{hid}$$

**Higgs mass tells us where  
to look for axion!**

$$m_a \sim 6.2 \mu\text{eV} \left( \frac{10^{12} \text{ GeV}}{f_a} \right)$$

# Little Hierarchy from radiative PQ breaking? exhibited within context of MSY model

Murayama, Suzuki, Yanagida (1992);  
Gherghetta, Kane (1995)

Choi, Chun, Kim (1996)


Bae, HB, Serce, PRD91 (2015) 015003

augment MSSM with PQ charges/fields:

$$\hat{f}' = \frac{1}{2} h_{ij} \hat{X} \hat{N}_i^c \hat{N}_j^c + \frac{f}{M_P} \hat{X}^3 \hat{Y} + \frac{g}{M_P} \hat{X} \hat{Y} \hat{H}_u \hat{H}_d.$$

$$M_{N_i^c} = v_X h_i |_{Q=v_X}$$

$$\mu = g \frac{v_X v_Y}{M_P}.$$





Large  $m_{3/2}$  generates small  $\mu \sim 100 - 200$  GeV!

# Why do soft terms take on values needed for natural (barely-broken) EWSB? string theory landscape?

- assume model like MSY/CCK where  $\mu \sim 100$  GeV
- then  $m(\text{weak})^2 \sim |m_{H_u}^2|$
- If all values of SUSY breaking field  $\langle F_X \rangle$  equally likely, then mild (linear) statistical draw towards large soft terms
- This is balanced by anthropic requirement of weak scale  $m_{\text{weak}} \sim 100$  GeV

*Anthropic selection of  $m_{\text{weak}} \sim 100$  GeV:*  
If  $m_W$  too large, then weak interactions  $\sim (1/m_W^4)$  too weak  
weak decays, fusion reactions suppressed  
elements not as we know them






statistical draw to large soft terms balanced by anthropic draw toward red ( $m(\text{weak}) \sim 100$  GeV): then  $m(\text{Higgs}) \sim 125$  GeV and natural SUSY spectrum!

Giudice, Rattazzi, 2006

HB, Barger, Savoy, Serce, PLB758 (2016) 113





statistical/anthropic draw toward FP-like region



## What happens to SUSY WIMP dark matter?

- higgsino-like WIMPs thermally underproduced
- 3 not four light pions  $\Rightarrow$  QCD theta vacuum
- EDM(neutron)  $\Rightarrow$  axions: no fine-tuning in QCD sector
- SUSY context: axion superfield, axinos and saxions
- DM= axion+higgsino-like WIMP admixture
- DFSZ SUSY axion: solves mu problem with  $\mu \ll m_{3/2}$ !
- ultimately detect both WIMP and axion!

usual picture

=>

mixed axion/WIMP




KJ Bae, HB, Lessa, Serce

much of parameter space is axion-dominated  
with 10-15% WIMPs




$\Rightarrow$





higgsino abundance

axion abundance




mainly axion CDM  
 for  $f_a < \sim 10^{12}$  GeV;  
 for higher  $f_a$ , then  
 get increasing wimp  
 abundance

# Direct higgsino detection rescaled for minimal local abundance

Bae, HB, Barger, Savoy, Serce


$$\mathcal{L} \ni -X_{11}^h \bar{\tilde{Z}}_1 \tilde{Z}_1 h$$

$$X_{11}^h = -\frac{1}{2} (v_2^{(1)} \sin \alpha - v_1^{(1)} \cos \alpha) (g v_3^{(1)} - g' v_4^{(1)})$$



Deployment of Xe-1ton,  
LZ, SuperCDMS  
coming soon!

Can test completely with ton scale detector  
or equivalent (subject to minor caveats)



range of  $f_a$  expected from SUSY  
with radiatively-driven naturalness  
compared to ADMX axion reach

# Conclusion: SUSY IS alive and well!

- old calculations of naturalness over-estimate fine-tuning
- naturalness: Little Hierarchy  $\mu \ll m(\text{SUSY})$  allowed
- radiatively-driven naturalness:  $\mu \sim 100\text{--}200$  GeV,  $m(t_1) < 3$  TeV,  $m(\text{gluino}) < 4$  TeV
- SUSY DFSZ axion: solve strong CP, solve SUSY  $\mu$  problem; generate  $\mu \ll m(\text{SUSY})$
- landscape pull on soft terms towards RNS,  $m(h) \sim 125$  GeV
- natural NUHM2: HL-LHC can cover via  $SSdB + Z1Z2j$  channels (see talk by X. Tata)
- expect ILC as higgsino factory
- DM = axion+higgsino-like WIMP admixture: detect both?





# Prospects for discovering SUSY

with radiatively-driven naturalness  
at LHC and ILC



# gluino pair cascade decay signatures

NUHM2:  $m_0=5 \text{ TeV}$ ,  $A_0=-1.6m_0$ ,  $\tan\beta=15$ ,  $\mu=150 \text{ GeV}$ ,  $m_A=1 \text{ TeV}$



Particle	dom. mode	BF
$\tilde{g}$	$\tilde{t}_1 t$	$\sim 100\%$
$\tilde{t}_1$	$b \tilde{W}_1$	$\sim 50\%$
$\tilde{Z}_2$	$\tilde{Z}_1 f \bar{f}$	$\sim 100\%$
$\tilde{Z}_3$	$\tilde{W}_1^\pm W^\mp$	$\sim 50\%$
$\tilde{Z}_4$	$\tilde{W}_1^\pm W^\mp$	$\sim 50\%$
$\tilde{W}_1$	$\tilde{Z}_1 f \bar{f}'$	$\sim 100\%$
$\tilde{W}_2$	$\tilde{Z}_i W$	$\sim 50\%$


Table 1: Dominant branching fractions of various sparticles along the RNS model line for  $m_{1/2} = 1 \text{ TeV}$ .

Int. lum. ( $\text{fb}^{-1}$ )	$\tilde{g}\tilde{g}$
10	1.4
100	1.6
300	1.7
1000	1.9

LHC14 5sigma reach  
in  $m(\text{gluino})$  (TeV)



since  $m(\text{gluino})$  extends to  $\sim 4 \text{ TeV}$ ,  
LHC14 can see about half the low EWFT  
parameter space in these modes

LHC14 has some reach for  
gluino pair production in RNS;  
if a signal is seen,  
should be distinctive




OS/SF dilepton mass  
edge apparent from  
cascade decays  
with  $z_2 \rightarrow z_1 + l + l^{\text{bar}}$

# Distinctive same-sign diboson (SSdB) signature from SUSY models with light higgsinos!




This channel offers best reach of LHC14 for RNS;  
it is also indicative of wino-pair prod'n  
followed by decay to higgsinos

Good old  $m_0$  vs.  $m_{1/2}$  plane still viable, but needs  $\mu \sim 100\text{--}200$  GeV as possible in NUHM2 instead of CMSSM/mSUGRA




See direct higgsino pair production recoiling from ISR (monojet signal)?



typically 1% S/BG after cuts:  
very tough to do!




What about  $pp \rightarrow \tilde{Z}_1 \tilde{Z}_2 j$  with  $\tilde{Z}_2 \rightarrow \tilde{Z}_1 \ell^+ \ell^-$  ?

Han, Kribs, Martin, Menon, PRD89 (2014) 075007;  
HB, Mustafayev, Tata, PRD90 (2014) 115007;







use MET to construct  $m^2(\text{tau-tau})$



# LHC reach for soft dilepton+jet+MET




# panoramic view of reach of HL-LHC for natural SUSY



LHC14 with  $3000 \text{ fb}^1$  can cover essentially all parameter space with  $\Delta_{EW} < 30$ , usually with 2-3 distinct signals:  $\tilde{g}\tilde{g}$ , SSdB and  $\tilde{Z}_1\tilde{Z}_2j$

# Smoking gun signature: light higgsinos at ILC:

ILC is Higgs/higgsino factory!




$$\sigma(\text{higgsino}) \gg \sigma(Zh)$$

10–15 GeV higgsino mass gaps no problem in clean ILC environment

HB, Barger, Mickelson, Mustafayev, Tata  
arXiv:1404.7510

ILC either sees light higgsinos or MSSM dead

# Future collider reach for naturalness




Bae, HB, Nagata, Serce

When to give up on naturalness in MSSM?  
 If HL-LHC or ILC(600GeV) sees no light higgsinos; WIMP at Xe-1ton/LZ

# Conclusions: status of SUSY post LHC8


- SUSY EWFT **non-crisis**: EWFT allowed at 10% level in radiatively-driven natural SUSY: SUGRA GUT paradigm is just fine in NUHM2 but CMSSM/others fine-tuned
- naturalness maintained for  $\mu \sim 100\text{--}200$  GeV;  $t_1 \sim 1\text{--}3$  TeV,  $t_2 \sim 3\text{--}8$  TeV, highly mixed;  $m(\tilde{g}, \tilde{t}) \sim 1\text{--}4$  TeV
- LHC14 w/  $3000 \text{ fb}^{-1}$  can see all  $\text{DEW} < 30$  RNS parameter space
- **$e^+e^-$  collider with  $\sqrt{s} \sim 500\text{--}600$  GeV needed to find predicted light higgsino states**
- Discovery of and precision measurements of light higgsinos at ILC!
- SUSY DFSZ/MSY invisible axion model: solves strong CP and SUSY  $\mu$  problems while allowing for  $\mu \sim m(Z) \ll m(\text{SUSY})$
- soft terms pulled to natural SUSY/barely broken EWS values, landscape?
- RNS spectra characterized by mainly higgsino-like WIMP: standard relic underabundance
- Expect mainly axion CDM with 5–10% higgsino-like WIMPs over much of p-space
- Ultimately detect **both axion and higgsino-like WIMP**



# Dark matter in RNS



Mainly higgsino-like WIMPs thermally underproduce DM



green: excluded;  
red/blue: allowed

HB, Barger, Mickelson

Factor of 10–15 too low

But so far we have addressed only **Part 1**  
of fine-tuning problem:

In QCD sector, the term  $\frac{\bar{\theta}}{32\pi^2} F_{A\mu\nu} \tilde{F}_A^{\mu\nu}$  must occur

But neutron EDM says it is not there: strong CP problem

(frequently ignored by SUSY types)

Best solution after 35 years:

PQWW/KSVZ/DFSZ **invisible axion**

In SUSY, axion accompanied by axino and saxion

Changes DM calculus:

expect mixed WIMP/axion DM (**2 particles**)

## Axion cosmology

★ Axion field eq'n of motion:  $\theta = a(x)/f_a$

$$- \ddot{\theta} + 3H(T)\dot{\theta} + \frac{1}{f_a^2} \frac{\partial V(\theta)}{\partial \theta} = 0$$

$$- V(\theta) = m_a^2(T) f_a^2 (1 - \cos \theta)$$

– Solution for  $T$  large,  $m_a(T) \sim 0$ :

$$\theta = \text{const.}$$

–  $m_a(T)$  turn-on  $\sim 1$  GeV


★  $a(x)$  oscillates,

creates axions with  $\vec{p} \sim 0$ :

production via vacuum mis-alignment

$$\star \Omega_a h^2 \sim \frac{1}{2} \left[ \frac{6 \times 10^{-6} \text{ eV}}{m_a} \right]^{7/6} \theta_i^2 h^2$$


★ astro bound: stellar cooling  $\Rightarrow f_a \gtrsim 10^9 \text{ GeV}$




# mixed axion-neutralino production in early universe

- neutralinos: thermally produced (TP) or NTP via  $\tilde{a}$ ,  $s$  or  $\tilde{G}$  decays
  - re-annihilation at  $T_D^{s,\tilde{a}}$
- axions: TP, NTP via  $s \rightarrow aa$ , bose coherent motion (BCM)
- saxions: TP or via BCM
  - $s \rightarrow gg$ : entropy dilution
  - $s \rightarrow SUSY$ : augment neutralinos
  - $s \rightarrow aa$ : dark radiation ( $\Delta N_{eff} < 1.6$ )
- axinos: TP
  - $\tilde{a} \rightarrow SUSY$  augments neutralinos
- gravitinos: TP, decay to SUSY


# DM production in SUSY DFSZ: solve eight coupled Boltzmann equations






higgsino abundance

axion abundance



mainly axion CDM  
 for  $f_a < \sim 10^{12}$  GeV;  
 for higher  $f_a$ , then  
 get increasing wimp  
 abundance

# Higgsino detection via halo annihilations:



green: excluded by Xe-100

annihilation rate is high but rescaling is **squared**

Gamma-ray sky signal is factor 10-20 below current limits